

**VISCOMETIC, VOLUMETRIC AND ULTRASONIC VELOCITY STUDIES OF
BINARY MIXTURES OF 2- PROPANOL AND 1-HEPTANOL WITH O-
NITROTOLUNE AT 303.15 AND 313.15K.**

R.D. Pawar¹ S.R. Patil² G.P. Waghulde³.

1- Arts, Science and Commerce, College, Yawal , (Jalgaon), M.S. India.

2- Arts, Science and Commerce, College, Chopda , (Jalgaon), M.S. India.

3- D.D.N. Bhole College Bhusawal (M.S)

Abstract :

The physicochemical properties like ultrasonic velocities, viscosities and densities of binary liquid mixtures of 2-propanol and 1-heptanol with o-nitrotolune were reported at 303.15 and 313.15 K. The Excess molar volume (V^E), Isentropic compressibility(ΔK_s) and viscosity deviation ($\Delta \eta$) have been calculated. These values were fitted with Redlich -Kister type polynomial equation. The results were interpreted in terms of molecular interaction between the components of the mixtures.

Keyword: Ultrasonic velocity, viscosity, density, Excess molar volume (V^E), Isentropic .

Introduction

The viscosity ,Density and Ultrasonic velocities measured find various application in characterising the physico- chemical properties of liquid mixtures ¹⁻³ and the study of molecular interaction . The ultrasonic velocity of liquid is related to the binding forces between atoms in the molecules .Ultrasonic velocity has been also employed in understanding the nature of molecular interactions in pure liquid ⁴and binary mixtures .The method studying the molecular interaction from the knowledge of variation thermodynamics parameters and their excess value with composition gives an insight into the molecular process ⁵⁻⁷ . The investigation regarding the molecular association in organic binary mixtures having one alkanol group as one of the components is of particular interest since 1- alkanol is highly polar and can associate with any other group having some degree of polar attraction. O-nitrotolune is strongly associated due to highly polar N=O group.

In view of the importance mentioned , an attempt has been made to elucidate the molecular interactions the mixture of O-nitrotolune with 2- propanol and 1 - heptanol respectively at 303.15 and 315.15 K further in the excess values of some of associated ultrasonic velocity , density and viscosity of mixture

Materials and Methods:

The chemicals O-nitrotolune with 2- propanol and 1-heptanol used were of analytical grade (A.R) minimum assay of 99.9% obtained from s. d. fine chemicals India. Which are used as such without further purification. The densities of pure components and binary mixtures were measured by using a Bi-capillary pycnometer. The purities of the above chemicals were checked density determination. The binary liquid mixtures of different known concentration were prepared in stopper measuring flask. The weight of the sample was measured using electronic digital balance with an accuracy of ± 0.1 mg. The viscosity was measured using gubbelohde viscometer (20ml) and the efflux time was determined using a digital clock to within ± 0.015 .



The ultrasonic velocity (U) in liquid mixtures have been measured using an ultrasonic interferometer (Mittal type, model F-81) working at 2 MHz frequency. The accuracy of sound velocity was $\pm 0.1 \text{ ms}^{-1}$.

Theory and Calculations:

Excess volumes of the mixtures have been calculated using density and mole fraction data given by equation:

$$V^E = (M_1X_1 + M_2X_2) / \rho_{12} - (M_1X_1) / \rho_1 - (M_2X_2) / \rho_2 \quad \text{--(1)}$$

Viscosity of Binary Mixtures is calculated by:

$$\ln \eta_m = X_1 \ln \eta_1 + X_2 \ln \eta_2 \quad \text{--(2)}$$

The measured viscosities of the mixtures have been used to obtain deviation in Viscosity parameters on the basis of linearity in following way,

Deviation in Viscosity of Binary Mixtures is calculated by :

$$\Delta \eta_m = \eta_{12} - X_1 \eta_1 - X_2 \eta_2 \quad \text{--(3)}$$

Deviation in isentropic compressibility have been evaluated by using the equation

$$\Delta k_s = k_s - (\Phi_1 k_{s1} + \Phi_2 k_{s2}) \quad \text{--(4)}$$

where k_{s1} , k_{s2} and k_s are isentropic compressibility of liquid mixtures and Φ is volume fraction of pure i^{th} component in the mixture and is defined as

$$\phi = (x_i V_i) / (\sum x_i V_i) \quad \text{--(5)}$$

where x_i and V_i are mole fraction and molar volume of i^{th} component in the mixture

Results and Discussion

In pure state, the self association of alkanols decreases with increasing chain length, when alkanols mixed with *o*-nitrotoluene then there is interaction between their individual functional groups (-OH and -NO₂). The presence of electron withdrawing group on benzene ring decreases electron densities. The polarity of alkanols is less hence there is degree of self association is less as compared to nitrotoluene⁹.

The experimental values of density, viscosity and ultrasonic velocity and value of excess volume, viscosity deviation and deviation in isentropic compatibility parameter for the two binary liquid system at 303.15 and 313.15 are given table 1 and 2.

The excess volume and viscosity deviation are negative over the entire mole fraction of alkanol 303.15 and 313.15 K temperature.

The excess of parameter of particular mole fraction of alkanols becomes less negative with increases of temperature. The negative value may be attributed to existence of dispersion and dipolar forces between unlike molecules and related to the differences in size and shape of molecules¹⁰.

The magnitude of $\Delta \eta$ and Δk_s , the sign and the extent of deviation of these properties from ideality depends on the strength of interaction between unlike molecules. According to Fort et.al. the excess viscosity gives the strength of the molar interaction between in molecules.

It is found that for the solution a good agreement was found in between Redlich-Kister parameters the solution of the fifth degree polynomial obtained with V^E , $\Delta \eta$ and Δk_s .

The measurement of viscosity in binary mixture yield some reliable in the study of molecular interaction from the given table it shows that the value of excess viscosity and deviation in isentropic compressibility decreases with increases in concentration of alkanols but however it found to increases with elevation of temperature.



Conclusion

The experimental data of ultrasonic velocity, density and viscosity are reported by binary mixture of 2- propanol and 1- heptanol with 0-nitrotoluene over entire range of mole fraction at 303.15 and 313.15 K calculated viscosity deviation , excess molar volume and the change with isentropic compressibility are fitted with Redlich –Kister type polynomial equation . Very large negative deviation are observed for the both the investigated system. This reveals the existence of molecular interaction in the binary mixtures . The present investigation shown that greater molar interaction exist in binary mixtures.

Table.1. Values of densities,viscosities,ultrasonic velocity,Excess molar volumes and Deviation in viscosity and deviation in isentropic compressibility for binary system of 2-propanol and O-nitrotoluene at 303.15 and 313.15 K.

Temp K	X ₁	P (gm/cm ³)	$\square \times 10^3$ (Nsm ⁻²)	U (M S ⁻¹)	V ^E × 10 ⁶ (m ³ /mole)	$\Delta \square \times 10^3$ (Kg m ⁻¹ s ⁻¹)	$\Delta k_s \times 10^{11}$ (m ² N ⁻¹)
	0.0000	0.77910	1.79040	1361.6	0.0000	0.000	0.00
	0.0466	0.80580	1.63390	1383.6	-0.5112	-15.481	-13.09
	0.0991	0.83140	1.48030	1393.2	-0.7948	-30.630	-8.82
	0.1583	0.86370	1.43900	1433.8	-1.6218	-34.522	-30.79
	0.2270	0.89000	1.36380	1484.0	-1.6993	-41.765	-46.68
303.15	0.3049	0.92680	1.36190	1598.5	-2.7114	-41.642	-95.98
	0.3972	0.96420	1.34260	1642.6	-3.6042	-43.201	-92.18
	0.5058	1.01510	1.33430	1655.5	-5.6897	-43.595	-73.01
	0.6375	1.02490	1.32830	1663.8	-3.5259	-43.665	-32.50
	0.7980	1.04430	1.31450	1692.1	-1.9550	-44.400	-0.06
	1.0000	1.06470	1.75020	1831.2	0.0000	0.000	0.00
	0.0000	0.77060	1.40180	1316.1	0.0000	0.000	0.00
	0.0466	0.79680	1.29130	1348.8	-0.0461	-10.639	-28.66
	0.0991	0.82240	1.20480	1383.3	0.1460	-18.809	-50.17
	0.1583	0.85450	1.17870	1450.9	-0.1340	-20.879	-94.65
	0.2270	0.88070	1.14140	1454.3	0.4319	-23.982	-75.60
313.15	0.3049	0.91700	1.13770	1466.3	0.1706	-23.640	-64.95
	0.3972	0.95660	1.13700	1533.3	-0.1007	-22.868	-83.43
	0.5058	1.00500	1.12560	1552.4	-0.9664	-23.016	-67.78
	0.6375	1.01360	1.11620	1567.9	2.6008	-22.754	-27.40
	0.7980	1.04450	1.10400	1618.6	4.4192	-22.508	-7.02
	1.0000	1.13740	1.31050	1682.4	0.0000	0.000	0.00

Table.2. Values of densities,viscosities,ultrasonic velocity,Excess molar volumes and Deviation in viscosity and deviation in isentropic compressibility for binary system of 1-heptanol and O-nitrotoluene at 303.15 and 313.15 K.

Temp K	X ₁	P (gm/cm ³)	$\square \times 10^3$ (Nsm ⁻²)	U (M S ⁻¹)	V ^E × 10 ⁶ (m ³ /mole)	$\Delta \square \times 10^3$ (Kg m ⁻¹ s ⁻¹)	$\Delta k_s \times 10^{11}$ (m ² N ⁻¹)
	0.0000	0.81720	4.96600	1553.2	0.0000	0.000	0.00
	0.0863	0.84900	4.29940	1596.9	-2.0413	-38.957	-33.24



	0.1750	0.86750	3.71500	1598.8	-1.6770	-68.873	-31.47
	0.2662	0.89560	3.25850	1610.3	-2.6587	-85.195	-38.54
	0.3609	0.93120	2.81730	1620.9	-4.4593	-98.862	-46.35
303.15	0.4585	0.95240	2.54040	1662.9	-3.9656	-95.165	-60.63
	0.5599	1.01060	1.87810	1665.4	-8.1137	-128.787	-67.93
	0.6640	1.07640	2.18920	1679.3	-12.4328	-64.201	-78.89
	0.7716	1.08150	1.89450	1705.4	-9.4805	-59.069	-73.11
	0.8841	1.09250	1.83760	1710.4	-7.0519	-28.581	-59.66
	1.0000	1.06470	1.75020	1631.2	0.0000	0.000	0.00
	0.0000	0.81040	3.78360	1533.1	0.0000	0.000	0.00
	0.0863	0.84170	3.35190	1583.0	-1.2156	-21.865	-35.17
	0.1750	0.86000	3.90990	1593.8	-0.0159	55.872	-34.84
	0.2662	0.88760	2.60070	1597.0	-0.1176	-52.494	-33.19
	0.3609	0.92290	2.33380	1601.2	-1.0560	-55.764	-33.40
313.15	0.4585	0.94410	2.15370	1610.7	0.3236	-49.636	-27.62
	0.5599	1.00190	1.57640	1617.1	-2.9310	-82.289	-32.56
	0.6640	1.06690	1.86020	1640.1	-6.2920	-28.164	-42.77
	0.7716	1.07210	1.68460	1646.5	-2.3289	-19.114	-22.44
	0.8841	1.08320	1.58260	1650.7	1.1456	-1.491	-0.79
	1.0000	1.13740	1.31050	1682.4	0.0000	0.000	0.00

REFERENCES :-

- 1 Kinocid, *J. Am. Chem. Soc.* S1,2950(1929).
- 2 K.S.Mehara, *Indian J. Pure and Appl. Phys.*, 38,760(2000).
- 3 R.J.Fortand W.R.Moore, *Trans Faraday Society*, 61,2102(1965)
- 4 Eyring, H.& Kincaid, J.F.(1938). Free volumes and free angles ratios of molecules in liquids. *J. Chem. Phys.*, 6,220-229
- 5 Mehta, S.K.& Chauhan, R.K.(1996). Ultrasonic velocity and apparent isentropic compressibilities in mixtures of non-electrolytes. *J. Sol. Chem.*, 26,295-308
- 6 Dewan, R.K., Gupta, C.M. & Mehta, S.K.(1988). Ultrasonic study of ethyl benzene + n-alkanols. *Acoustica*, 65,245.
- 7 Granberg L. and Nissan, *Nature*, 1946, 164, 799
- 8 Patil S.R., Deshpande. U. Gand Hiray. A. R. Rasayan *J. Chem*, 2010, 66-73.
- 9 Wankhede, D.S.; Lande, M.K.; Arbad, B.R. Densities & viscosities of Binary Mixtures of Paraldehyde + Propylene Carbonate at (288.15, 293.15, 298.15, 303.15 & 308.15) K. *J. Chem. Eng. Data* 2005, 50, 261-263.
- 10 Daakua, V.K.; Sinha, B.; Roy, M.N. Thermophysical Properties of N-N-Dimethylformamide with isometric butanols at 298.15, 308.15 & 318.15 K. *J. Ind. Chem. Soc.* 2007, 24, 37-45.
- 11 Alvarez, E.; Cancela, A.; Maceiras, R.; Navaza, J.M.; Taboas, R.; Density, Viscosity, Excess molar volume, and Viscosity Deviations of Three amyl Alcohols + Ethanol Binary Mixtures from 293.15 to 323.15 K. *J. Chem. Eng. Data* 2006, 51, 940-945.
- 12 S.B. Kasare and B.A. Patdai, *Indian J. Pure & Appl. Phys* 25 180 (1987)
- 13 S. Jyostna, S. Nallani, *J. Chem. Thermodyn.*, 38(3), 272 (2006), DOI:10.1016/j.jct.2005.05.06
- 14 A. Ali. A. K. Nain, *Indian J. Phys* 74B, 63(2000).
- 15 M. Dzida, *J. Chem Eng Data*, 52(2) 521 (2007).
- 16 Pramod P Patil, Rupali S Patil, S.R. Patil, Amulrao U Borse *Rasayan J. Chem.* 2018., 1103-1112