

# EFFECT OF COPPER SULPHATE ON GLYCOGEN CONTENT OF FRESH WATER BIVALVE *L. marginalis*.

✓ Sandhya M. Sonawane, Mayur M. Sonawane.

Nutan Maratha College, Jalgaon, Dist. Jalgaon.

E-mail I.D. : [sandhyamahendra11@gmail.com](mailto:sandhyamahendra11@gmail.com),

Mulji Jaitha College Jalgaon, Dist Jalgaon.

E-mail I.D. : [mayursonawane13@gmail.com](mailto:mayursonawane13@gmail.com),

## ABSTRACT

Presence of heavy metals in stream & rivers is largely due to the runoff from agricultural fields and outfall from manufacturing factories. Heavy metals are not highly selective but are generally toxic to many non-target organisms. The aquatic environment is also polluted by heavy metals & leads to many changes in organism physiology. In the present investigation an attempt has been made to observe the effect of Copper sulphate on Glycogen content of *Lamellidens marginalis*. After acute & chronic treatment with Copper sulphate the total Glycogen content of digestive gland of bivalve was altered significantly than foot, whole body and mantle ( $P < 0.001$ ), indicating altered carbohydrate metabolism.

## INTRODUCTION

Most information about the effect of environmental pollutants on aquatic animals has been obtained from mortality studies. Often very little is known about damage to different internal organs or about disturbed physiological and biochemical processes within an organism following exposure to environmental poisons. Consequently knowledge about the mode of action of toxicants and causes of death in poisoned aquatic animals is often lacking. A better understanding of these mechanisms is necessary if we want to predict the potential harmfulness of various chemicals to the environment.

Since different environment pollutants are likely to affect biological systems in different ways according to their respective chemical properties, the sum of physiological changes created by a particular pollutant is likely to be characteristic of that pollutant. Thus by observing the effects of pollutants on a set of physiological parameters, it might be possible to establish specific responses of that pollutant, and may take it possible to identify a pollutant on the basis of its physiological effect pattern.

The higher concentrations of toxicants bring the adverse effects on aquatic organisms, at cellular level or molecular level and ultimately lead to disorder in biochemical composition which is useful in determining different toxicants and protective mechanism of the body to resist the toxic effects of the substances.

The toxic chemicals (pollutants) act as one kind of stress to organism and organism responds to it by developing necessary potential to counter act that stress. The biochemical changes occurring act that stress. The biochemical changes occurring in the body give first indication of stress. During stress the organism needs sufficient energy which is supplied from reserve materials (glycogen, lipid and protein). If the stress is mild then only stored glycogen is used, as a source of energy, but if the stress is strong then the energy stored in lipid and protein may be used.

The heavy metals cause metabolic derangement in the living system, when come in contact. Heavy metals due to their potential toxicity produce biochemical changes in the tissues of animals.

Even though there is much work on the toxic effects of pesticides and heavy metals on specific target animals, little attention was paid to study the physiological and

biochemical changes on non target aquatic species. Since *Lamellidens Marginalis* are fresh water bivalves an attempt was made to study the changes in biochemical composition by heavy metals copper sulphate. Carbohydrates (glycogen) constitute the vital organic substances playing an important role in energy metabolism. So a change in total glycogen is studied in the present investigation.

The biochemical constituents were estimated from whole body, foot, digestive gland and mantle. These tissues are vital and metabolic important and any stress on the animal is depicted by the changes in the constituent in these tissues. Carbohydrate plays structural role in every living organism and serves as a reservoir of the chemical energy and many decrease or increase according to needs of the organism. The tissue glycogen is a major source for energy for metabolic processes (Lather, 1975).

A change in glycogen content in various organisms after exposure to various pollutants was studied by many workers (Peter, 1973; Mayers, 1977; Koundinya and Ramamurthi, 1979; Bhagyalakshmi, 1981; Gill and Pant 1981; Patil 1986; and Choudhari, 1988). Baner and Ghosh (1978) has recorded a change in levels of serum glucose, liver glycogen and glucose-6-phosphate of fishes, *Clarius batrachus* and *Tilapia mossambica* when exposed to cadmium. Koundinya and Ramamurthi (1978) observed the effect of lethal concentration of sumithion on carbohydrate metabolism of the Indian cat fish, effect of methyl parathion on carbohydrate metabolism of the Indian cat fish, *Heteropneustes fossilis* was determined by Srivastava and Singh (1981). The effect of various concentrations of zinc sulphate on glycogen content of liver, muscle, brain, kidney and grills of air breathing freshwater food fish, *Anabas Scandus* has been investigated by Natarajan (1982).

A little information is available about biochemical diversions due to heavy metals and pesticide pollutant stress in Molluscs. Mandal and Ghose (1970) studied the effect of cadmium arsenate on the snail *Achatina Fulica* has (Bawdich) on mobilization of digestive gland glycogen. Ramanarao and Ramamurthi (1980) studied the effect of sublethal concentration of sumethion on biochemical composition of the snail, *Pila globosa*. Chaudhari (1988) studied the effect of pesticides on the freshwater prosobranch snail, *Bellamya bengalensis* with respect to change in glycogen.

## MATERIAL AND METHODS

The bivalves, *Lamellidens marginalis* were collected from the Godavari river at Paithan. They were brought to the laboratory and maintained in plastic troughs. The bivalves were acclimatized to laboratory conditions for two-three days. The water in the troughs was changed every day. The healthy adult bivalves of approximately the same size and weight were selected for the experiment. During the chronic treatment the animals were fed on algae or *Hydrilla*.

To study the effects of heavy metals (copper sulphate) on the biochemical composition of *Lamellidens marginalis*, the bivalves were exposed to median lethal concentration and sub lethal concentration of heavy metals as acute treatment and chronic treatment respectively.

### (A) Acute treatment :

The acclimatized bivalves were divided into 2 groups. The first group was exposed to 1.6 ppm copper sulphate for 72 hours. The second group of bivalve was kept as control.

At the end of 24, 48 and 72 hours treatment on pollutant copper sulphate, the control and treated bivalves were scarified to analyze the biochemical composition. The bivalves were dissected and their foot, digestive gland and mantle were separated. Then glycogen contents in the tissues of treated and control bivalves were analyzed. The percentages of biochemical components in the tissues of treated and control bivalves were compared.

### (B) Chronic treatment:



The acclimatized bivalves were divided into two groups, one group of bivalves was kept as control and other group of bivalves was exposed to 0.82 ppm copper sulphate. The chronic treatment was given up to 20 days. The control and treated bivalves were fed on freshwater algae or Hydrilla during exposure period.

At the end of the 5<sup>th</sup>, 10<sup>th</sup>, 15<sup>th</sup> and 20<sup>th</sup> day, the control and treated bivalves were sacrificed to analyze the biochemical composition. The control and treated bivalves were dissected and their foot, digestive gland and mantle were separated. The glycogen contents in the tissues (whole body, foot, digestive gland and mantle) of treated and control bivalves were analyzed. The percentages of biochemical components in the tissues of control and treated bivalves were compared.

#### **Estimation of Biochemical components from tissues : (Glycogen)**

The colorimetric estimation of glycogen present in the tissue was done by anthrone reagent method (Dezwaan and Zandee, 1972). 50 mg. of wet tissue was taken in 1 ml of 30% KOH solution. The mixture was boiled in water bath for 5-10 minutes, till the tissue was completely dissolved. The solution was cooled and to it 0.2 ml 2% Na<sub>2</sub>SO<sub>4</sub> and 6 ml of absolute alcohol were added. This solution was kept in refrigerator for overnight. It was then centrifuged for 15 minutes at 3000 rpm. The supernatant was discarded and the residue cake was dissolved in 10 ml of distilled water 0.1 ml of this solution was taken and to it 0.9 ml of distilled water were added. The solution was heated in boiling water bath for 5 minutes and then cooled. The intensity of the colour developed was measured with the colorimeter (Erma) at 620 mμ (Red Filter) filter. Anthrone, reagent was prepared by dissolving 50 mg anthrone powder and 1 gm Thiourea in 100 ml of 72% H<sub>2</sub>SO<sub>4</sub>. The amount of glycogen was calculated by referring to a standard graph value, where glucose was used as a standard. The glycogen value was calculated by multiplying with the conversion factor 0.927 to glucose value. The amount of glycogen was expressed in terms of Mg. of glycogen/100 mg of wet tissue.

#### **OBSERVATION AND RESULTS**

Biochemical components such as carbohydrates (glycogen) was studied in the normal (control) and pollutant (copper sulphate) treated whole body, foot, digestive gland and mantle of freshwater bivalve, *L. marginalis*., the results are summarized in Tables .

A) Acute treatment-The acute treatment was given upto 72 hours by heavy metal pollutant, copper sulphate. After the acute treatment by pollutant, biochemical composition of the bivalve was altered and the results are summarized in the tables .The glycogen content of the whole body, foot digestive gland and mantle decreased after acute treatment by heavy metals. After copper sulphate acute treatment at 72 hours, The glycogen in the whole body, foot, digestive gland and mantle decreased from 2.08% to 1.22%, 2.53% to 2.13%, 2.78% to 1.62% and 1.13% to 0.86% respectively. The glycogen content of the foot and mantle depleted more as compared to the digestive gland.

B) Chronic treatment-The biochemical components such as glycogen was observed after chronic treatment in control and treated freshwater mussels, *Lamellidens marginalis*. The biochemical components of the whole body foot, digestive gland and mantle were observed and recorded in the table. The glycogen content of the whole body, foot, digestive gland and mantle decreased in chronic (5, 10, 15 and 20 days) treatment by heavy metal. After copper sulphate chronic treatment, the glycogen in the whole body foot, digestive gland and mantle decreased..The maximum depletion occurred in the digestive gland of bivalve.

Table – 1

Glycogen content in selected tissues of the control and CuSO<sub>4</sub> exposed *Lamellidens marginalis* as a function of exposure period

Treatment	Sr. No.	Body Organ	Total glycogen content (%) + S.D.		
			24 hours	48 hours	72 hours
Control	1	Whole body	2.2874 ± 0.0713	2.1475 ± 0.002	2.0879 ± 0.0711
	2	Foot	2.8725 ± 0.00520	2.9675 ± 0.0578	2.5324 ± 0.0307
	3	Digestive gland	3.2035 ± 0.0036	3.0234 ± 0.0128	2.7852 ± 0.0787
	4	Mantle	1.2242 ± 0.0179	1.2542 ± 0.0427	1.1342 ± 0.0126
Acute treatment by CuSO <sub>4</sub>	1	Whole body	1.7529 ± 0.0438 P<0.01	1.6432 ± 0.0189 P<0.001	1.1272 ± 0.0234 P<0.001
	2	Foot	2.6435 ± 0.0374 NS	2.6720 ± 0.2203 NS	2.1382 ± 0.01156 P<0.05
	3	Digestive gland	2.2705 ± 0.1328 P<0.001	2.0035 ± 0.0857 P<0.001	1.6240 ± 0.1239 P<0.001
	4	Mantle	0.8900 ± 0.0179 P<0.001	0.8899 ± 0.0245 P<0.001	0.8601 ± 0.0554 P<0.001



Table – 2

Glycogen content in selected tissues of the control and CuSO<sub>4</sub> exposed *Lamellidens marginalis* as a function of exposure period

Treatment	Sr. No.	Body Organ	Total glycogen content (%) + S.D.			
			5 Days	10 Days	15 Days	20 Days
ontrol	1	Whole body	2.2378 ± 0.0243	2.1879 ± 0.0732	2.1004 ± 0.0002	1.8907 ± 0.0731
	2	Foot	2.2021 ± 0.0009	3.1357 ± 0.0306	2.3820 ± 0.0915	2.7692 ± 0.0531
	3	Digestive gland	3.92130 ± 0.0831	3.1242 ± 0.0101	2.8012 ± 0.0071	0.6402 ± 0.0432
	4	Mantle	1.3456 ± 0.0524	1.3000 ± 0.0861	1.0957 ± 0.0769	0.9802 ± 0.0216
Acute treatment by CuSO <sub>4</sub>	1	Whole body	1.2839 ± 0.0658 P<0.001	1.0897 ± 0.0723 P<0.001	0.9945 ± 0.0787 P<0.001	0.8445 ± 0.0307 P<0.001
	2	Foot	1.8403 ± 0.0581 P<0.001	1.5002 ± 0.0881 P<0.001	1.4640 ± 0.0357 P<0.001	1.0041 ± 0.0011 P<0.001
	3	Digestive gland	1.8432 ± 0.1254 P<0.001	1.5308 ± 0.1162 P<0.001	1.2920 ± 0.0762 P<0.001	1.0025 ± 0.0020 P<0.001
	4	Mantle	0.8280 ± 0.0199 P<0.001	0.6625 ± 0.0523 P<0.001	0.4324 ± 0.0134 P<0.001	0.3425 ± 0.015 P<0.001

## DISCUSSION

Heavy metal and pesticide pollutants cause stress to the aquatic organism and change its metabolic activity. The change in biochemical composition of tissues due to heavy metals and pesticides and the physiological state of metabolic activity of an organism reflect the utilization of their biochemical energy to counteract the toxic stress. The observed biochemical changes in bivalves representing adaptive or regulatory mechanism may be due to pathological effect. The animal by changing its metabolic processes tries to overcome the toxic effects as a protective measure. The heavy metal pollutants give rise to alterations in the metabolic and physiological activity both after short and long term exposures. To investigate the physiological changes after pesticide and heavy metal treatment the most fundamental one would be the study of change in the biochemical constituents. Carbohydrates, proteins and lipids are the important metabolites which provide energy to different vital processes.

Glycogen is the stored food material in animal tissue which is used as an immediate source of energy when required and is an essential feature of the normal organismal metabolism (Thunberg and Manchester, 1972.)

They glycogen content in freshwater bivalve, *L. marginalis* was altered indicating the effects of heavy metals. The average glycogen content in acute and chronic treatment by heavy metals (copper sulphate, mercury chloride, and cadmium chloride) was decreased in the whole body. The depletion of glycogen content was greater in the digestive gland as compared to the foot and mantle of the bivalve, when exposed to pollutants. This indicates that the digestive gland is the principal metabolic center for various metabolic functions. During acute and chronic exposures a significant decrease in the glycogen content of the digestive gland suggests greater glycolytic activity in the gland than the mantle and foot. The greater breakdown of glycogen may suggest the need of high energy to animal in stress conditions caused due to pollutants. Depletion in glycogen level might be because of the anoxia and hypoxia caused due to stress conditions which are known to increase carbohydrate consumption (Dezwan and Zandee, 1972).



The decrease in glycogen content indicates shifting towards anaerobic metabolism. Carbohydrates are considered to be the first organic nutrient to be depleted and degraded in stress conditions imposed on animals (Clerke, 1975). According to Koundinya and Ramamurthi (1979), the decrease in glycogen may be due to enhanced breakdown of glycogen to glucose through glycolysis. The greater decrease in the glycogen level, in the digestive gland might be due to high potential of digestive gland for glycolysis, similar to that of the vertebrate liver as suggested by Kabeer et al., (1977).

The mode of action of pollutants may be responsible for cellular dis-organization offering the storage and metabolism of the glycogen. Decrease in glycogen content indicates disrupted carbohydrate metabolism. The pollutants give the heavy physical irritative stress causing rapid movement and increased respiration rate thus increased utilization of reserved glycogen to meet higher energy demand of body causing decrease in glycogen content (Bhagyalaxmi, 1981). Many workers support the above results in vertebrate and invertebrate animals. A change in serum protein and glycogen content of rainbow trout exposed to endrin was studied by Grant and Mahrle (1973). Koundinya and Ramamurthi (1978) observed the effect of lethal concentration of sumethion on carbohydrate metabolism in *Tilapia mossambica*. Baner and Ghosh (1978) have recorded the alterations in the levels of serum glucose, liver glycogen and glucose-6-phosphate of the fish, *Clarius batrachus* when exposed to cadmium. Banerjee et al. (1978) reported an increase in blood glucose level in the fish *Tilapia mossambica* when exposed to cadmium. Kabeer (1979) stated that decrease in glycogen content in Malathion exposed tissues can also be due to decrease in glycogen synthesis. Koundinya and Ramamurthi (1979) have observed that sumethion leads to an increase in blood glucose level in decrease in glycogen content. Rao and Rao (1979) studied the effect of methyl parathion on the fish *Sarotherodon mossambica* and noted a significant decrease in glycogen content. Gill and Pant (1981) studied the effect of Nickel intoxication of carbohydrate metabolism i.e. blood glucose and liver glycogen which were measured to assess the magnitude of biochemical stress. Bhagyalakshmi (1981) studied the levels of certain carbohydrate metabolites in the tissue of field crab, *Oxiotelphusa senex senex* after exposure to an organophosphate pesticide sumithion, and recorded an increase in haemolymph glucose level and decrease in total carbohydrate level.

Nagabhushanam and Kulkarni (1981) observed increased haemolymph glucose and a decrease in midgut gland, when the prawns, *Macrobrachium Kistensis* were exposed to  $ZnSO_4$  and  $CuSO_4$ . Srivastava and Singh (1981) studied the acute effect of methyl parathion on carbohydrate metabolism of the Indian carfish, *Heteropneustes fossilis*. Forroqui (1982) observed a insignificant increase of glycogen in the ovary of *Barytelphusa cunicularis* after two days exposure to sevimol and a significant decrease after seven days exposure and concluded that glycogen breakdown provides the immediate energy source in stress condition. Natarajan (1982) studied the effect of various concentration of  $ZnSO_4$  on glycogen content of liver, muscle, brain, kidney and gills of the fish *Anabas Scandis*. Bhagyalakshmi et al. (1984) observed a decrease in glycogen and elevated phosphorylase activity in the crab, *Oxiotelphusa senex senex* exposed to sumithion, suggesting onset of glycogenolysis forming free glucose and the possible exist of these glucose molecules in the haemolymph resulting in the hyperglycemia condition. Patil (1986) studied the effects of pesticides on the glycogen content of *Mythimna (P)* separate and found decreased glycogen content after treatment.

Jalaluddin Shariff (1987) studied the acute toxicity of 1 det 10 L on biochemical composition of fish and found reduced glycogen content in the muscle after detergent treatment. Srinivasulu Reddy and Ramana Rao (1987) studied the effect of phosphamidon pesticide on glucose metabolism in different tissues (midgut, muscle and gill) and noted that the decrease in total carbohydrate levels signified their utility possibly to meet the higher energy demands and concluded that the both the lethal and sublethal concentrations of phosphamidon increase the oxidation of glucose through TCA cycle and enhance both glycolysis and hexose monophosphate pathways to yield more number of ATP and  $NADPH_2$  molecule for detoxification. Reddy et al. (1987) observed decreased glycogen level in the prawn, *Metapenaeus monoceros* after exposure to phosphamidon, DDT and Fevelrate. Vijaykumar and Kannupandi (1989) studied the effect of organophosphate insecticide phosphamidon on carbohydrate content and found decreased carbohydrate content in the larval stages of the crab, *Sesarma andersonia*.

Many workers studied the effects of pollutants on mollusks in Mandal and Ghose (1970) observed glycogen depletion in the digestive gland of the snail, *Achatina fulica* (Bawdich) when exposed to calcium arsenate. Ramana Rao and Ramamurthi (1980) studied the effect of sublethal concentration of sumithion on some biochemical constituents of the snail, *Pila globosa* and found a decrease in glycogen content. Lomte and Alam (1982) observed the stable level of glycogen in the foot and mantle but very sharp fall of glycogen in the digestive gland during sublethal exposure for 24



hours to organophosphate pesticide, malathion. Swami et al. (1983) found increased haemolymph glucose and decreased stored glycogen in *Lamellidens marginalis* when treated with Flodit and Metacid. Kulkarni et al. (1984) studied the impact of endosulfan on the apple snail, *Pila globosa* and found an elevation of blood glucose level after treatment. Chaudhari (1988) studied the effect of pesticides on biochemical composition of the snail *Bellamya (viiparous) bengalensis* and found decreased glycogen content after treatment.

## SUMMARY

1. The biochemical composition of *Lamellidens marginalis* after acute and chronic treatment of heavy metals (copper sulphate) was studied to better understand the mechanism of action of pollutant, by observing the time bound and tissue specific alterations of biochemical component glycogen, in different organs of the body.
2. After acute treatment of heavy metals glycogen content of the whole body, foot, digestive gland and mantle was altered very prominently, glycogen depletion was more due to copper sulphate as compared to control
3. After acute treatment of heavy metals ( $\text{CuSO}_4$ ,  $\text{HgCl}_2$  and  $\text{CdCl}_2$ ) the lipid content decreased in the whole body, foot and mantle of the bivalve, *L. marginalis*.
6. The glycogen content of the whole body, foot, digestive gland and mantle decreased after chronic treatment of heavy metal pollutants, much glycogen content decreased in digestive gland which was followed by mantle and foot. Glycogen decrease due to  $\text{CuSO}_4$ .

## REFERENCES

- Abel, P.D. (1974) : Toxicity of synthetic detergent to fish and aquatic vertebrates. *J. Fish. Biol.*, 6 : 279-298.
- Banerjee, S.K. and Ghosh, S.D. (1978) : Toxicity of cadmium : A comparative study in the air breathing fish *Clarius batrachus* (Linn.) and in the non air breathing one, *Tilapia mossambica* (Peters). *Ind. J. Exp. Biol.*, 16 : 1274.
- Banerjee, S.K., Dastidar, S.G., Mukhopadhyaya, P.K. and Dehadari, P.V. (1978) : Toxicity of cadmium : A comparative study in air breathing fish, *Clarius batrachus* (Linn.) and in non air breathing one *Tilapia mossambica* (Peters), *Indian J. Exp. Biol.*, 16 : 1274-1277.
- Calvin D.B. (1931) : Glycogen content of freshwater mussels. *Proc. Soc. Expt. Biol. Med.*, 29 : 96-97.
- Clarke, D.H. (1975) : *Exercises in Physiology*, I Parintice Hall, G. New Jersey.
- Choudhary, T.R. (1988) : Pesticidal impact on some physiological aspects of *Bellamya (viviparus) bengalensis*. Ph.D. thesis, Marathwada University, Aurangabad. M.S., India.
- Coley, R.M. and Jensen, R.B. (1973) : *Lipid*, 8 : 43.
- Coppage, D.L., Mathews, E., Cook, G.H. and Krught, J. (1975) : *Pest. Biochem. Physiol.*, 15 : 536.
- Copuzzo, J.M. and Lancaster, B.A. (1981) : Physiological effects of South Louisanna crude oil to the American Lobster (*Homarus americanus*) : Hydrocarbon accumulation interference with lipid metabolism. In physiological mechanism of marine pollutants toxicity, 477-507 (Eds.) W.B. Vernberg, A. Calabrese, F.P. Thurberg and F.J. Vernberg, New York, Academic Press.
- Daniel, R.J. (1922) : Seasonal changes in the composition of mussel (*Mytilus edulis*) Lancashire, Sea Fisheries Lab Report, 31 : 27-50.
- Dezwann, B. and D.I. Zandee (1972) : The utilization of glycogen and accumulation of some intermediates comp. *Biochem. Physiol.*, 43 B : 47-54.
- Durve V.S. and Bal D.V. (1961) : Studied on the chemical composition of the oyster, *Crossstrea gryphoides*. *J. Zool. Soc. Indian* 13 : 70-77.
- Cabbott, P.A. (1976) : Energy metabolism. In : Bayne B.L. (Ed.). *Marine mussels their ecology and physiology* I.B.P. Cambridge University Press, PP. 293-337.

- Holland, D.L. (1978) : Lipid reserves and energy metabolism. In larvae of benthic marine invertebrates : In Biochemical and Biophysical perspectives in marine biology, 85-123 (Eds.) D.C. Maline and J.R. Sargent, New York, Academic Press.
- Jalaluddin Shariff, D. (1987) : Modulation in the biochemical nature of the body muscles of a freshwater teleost *Sarotherodon mossambicus* (Peters) due to 1 Detol (A synthetic detergent). Proc. Nat. Cont. Env. Impact on biosystem. Pp. 279-282.
- Joseph, Thatheyus, A., M. Salvanayagam and T. Peter Raj (1987), Chronic toxicity of copper on the biochemical composition of some tissues of the scale carp, *Cyprinus carpio communis*. Proc. Nat. Con. Env. Impact on Biosystem. pp 263-267.
- Kabeer Ahmed, I. sivalah, S. and Ramana Rao, K.V. (1977) : On the possible significance of changes in organic constituents in selected tissues of Malathion exposed snail, *Pila globosa* (Swainson). Comp. Physiol. Ecol., 4(2) : 81-82.
- Kabeer Ahmed, I., Begum, M.R. Sivaiah, S. and Ramana Rao, K.V. (1978) : Effect of Malathion on free amino acids, total proteins, glycogen and some enzymes of pelecypod, *Lamellidens marginalis* (Lamark). Proc. Indian Acad. Sci. B. 87(12) : 377-380.
- Kabeer Ahmed Sahib, I. (1979) : studies on some aspects of protein metabolism and associated enzyme systems in the freshwater teleost *Tilapia mossambica* (Peters) subjected to Malathion exposure. Ph.D. Thesis, Sri Venkateswara University, Tirupati, A.P., India.
- Kaundinya P.R. and Ramamurthi P. (1979) : Effect of sublethal concentration of sumethion and sevin on certain haematological values of *Sarotherodon mossambicus*. Current Sciences, 49 : 645-646.
- Lather, L. (1975) : In chemical Biochemistry, W.B. Saunders Company, Philadelphia.
- Lomte, V.S. and Sabiha Alam, (1982) : Changes in the biochemical components of the prosobranch *Belamiya* (viviparous) *bengalensis* on exposure to malathion. Proc. Symp. Phys. Resp. Anim. Pollutants.
- Mandal T.K. and K.C.Ghose (1970) : Application of calcium arsenate : Histopathological changes and glycogen mobilization in *Achitina fulica* (Bawdich). Indian J. Exp. Biol., 8 : 332-333.
- Masumoto B., M. and Habino, M. (1934) : Biochemical studies of *Magaki* (*Ostrea gigas*) II. The seasonal variation in the chemical composition of *Strea gigas*. J. Sci. Hiros Hima Univ. A., 4 : 47-56.
- Muley, M.B. (1991) : Some physiological studies of *Melanoid tuberculatus* in relation to pollution stress. Ph.D. Thesis, Marathwada University, Aurangabad, M.S., India.
- Mitchell, P.H. (1915) : Nutrition of Oysters; glycogen formation and storage. Bull., U.S. Bur. Fish., 35 : 151-162.
- Nagabhushanam, R. and Kulkarni, G.K. (1981) : Freshwater *Palaemonid* Prawn, *Macrobrachium Kistensis* (Tiware) – Effect of heavy metal pollutants. Proc. Indian Metal Sci. Acad. 547 (3) : 380-386.
- Patil, P.N. (1986) : Impact of different pesticides on some physiological and neuroendocrinological aspects of *Mythima* (*Pseudaletia*) *separate*, Ph.D. Thesis, Marathwada University, Aurangabad, M.S., India.
- Peter, M. (1973) : Metabolism of Carbohydrates. In : Review of physiological chemistry, Ed. By Harper, H.A. Lange Medical Publication, 14<sup>th</sup> Edn, pp. 232-267.
- Ramana Rao, M.V. and Ramamurthi, R. (1978) : Studies on the metabolism of the apple snail, *Pila globosa* (Swainson in relation to pesticide impact). Ind. J. Her. 11 : 10.
- Ramana Rao, M.V. and Ramamurthi R. (1982) : Effect of exposure to sublethal concentration of mercuric chloride in some aspects of lipid metabolism in the fresh water field crab, *Oxiotelphusa senex senex* (F.) All India symposium on physiological responses of animals to pollutants, Marathwada University, Aurangabad, M.S., India.



- Rao, K.R., D.A. Kulkarni, K.S.Pillai and U.H.Mane (1987) : Effect of fluoride on the fresh water bivalve mollusks, *Indonaia caeruleus* (Prashad, 1918) in relation to the effect of pH biochemical approach. Proc. Nat. Symp. Ecotoxic, pp. 13-20.
- Sekeri, K.E., C.E. Sekeris, and P. Karlson, (1968) : Protein synthesis in Subcellular fractions of the blofly during different development stages. J. Insect. Physiol., 14 : 425-431.
- Thurberg, L.V. and Manchester, K.L. (1972) : Effect of dinervation on the glycogen content and on the activities of enzymes glucose and glycogen metabolism in rat diaphragm muscle. Biochem. J. 128 : 789.
- Ventakaraman, R. and Chary, S.T. (1951) : Studies on oysters and Clams, Biochemical Variations. Indi. J. Med. Res., 39(4) : 533-541.

